



Up-to-date Questions and Answers from authentic resources to improve knowledge and pass the exam at very first attempt. ----- Guaranteed.



NCEES PE Civil: Water Resources and Environmental Dumps  
NCEES PE Civil: Water Resources and Environmental Braindumps  
NCEES PE Civil: Water Resources and Environmental Real Questions  
NCEES PE Civil: Water Resources and Environmental Practice Test  
NCEES PE Civil: Water Resources and Environmental Actual Questions



[killexams.com](https://killexams.com)

**NCEES**

**NCEES PE Civil: Water Resources and Environmental**

*NCEES - PE Civil Engineering - Water Resources and Environmental 2024*

ORDER FULL VERSION

<https://killexams.com/pass4sure/exam-detail/NCEES-PE-Civil-WRE>



**Question 386:**

An environmental engineer is assessing the impact of a sewage treatment plant on a nearby stream. If the plant discharges effluent with a biochemical oxygen demand (BOD) of 200 mg/L and the stream's flow is 3 m<sup>3</sup>/s, what is the total BOD load entering the stream from the plant in kilograms per day?

- A. 51,840 kg/day
- B. 21,880 kg/day
- C. 45,320 kg/day
- D. 65,000 kg/day

**Answer:** A

**Explanation:** The BOD load can be calculated as:

$$\text{BOD Load} = \text{Concentration} \times \text{Flow Rate} \times \text{Time}$$

Convert mg/L to kg/m<sup>3</sup>:

$$\text{Concentration} = 200 \text{ mg/L} = 0.2 \text{ kg/m}^3$$

Thus,

$$\text{BOD Load} = 0.2 \text{ kg/m}^3 \times 3 \text{ m}^3/\text{s} \times 86,400 \text{ s} = 51,840 \text{ kg/day}$$

**Question 387:**

A groundwater engineer is evaluating the effects of a contaminant plume in a confined aquifer. If the hydraulic conductivity is 20 m/day and the contaminant concentration decreases from 1,000 µg/L to 100 µg/L over a distance of 50 m, what is the attenuation factor?

- A. 0.1

- B. 0.5
- C. 0.7
- D. 10.0

**Answer:** D

**Explanation:** The attenuation factor is calculated as:

$$\text{Attenuation Factor} = \frac{C_1}{C_2} = \frac{1000 \mu\text{g}/L}{100 \mu\text{g}/L} = 10$$

**Question 388:**

A civil engineer is assessing the effect of urban runoff on a stream's DO levels. If the stream's DO was 8 mg/L before the runoff event and dropped to 5 mg/L after, what is the percentage change in DO?

- A. 20%
- B. 25%
- C. 37%
- D. 35%

**Answer:** C

**Explanation:** The percentage change in DO is calculated as:

$$\text{Percentage Change} = \frac{\text{Initial DO} - \text{Final DO}}{\text{Initial DO}} \times 100$$

Thus,

$$\text{Percentage Change} = \frac{8 - 5}{8} \times 100 = 37.5\%$$

**Question 389:**

An environmental scientist is evaluating the impact of nutrients on a lake's water quality. If the lake has a volume of  $1,000,000 \text{ m}^3$  and the total phosphorus concentration is  $0.2 \text{ mg/L}$ , what is the total phosphorus load in kilograms?

- A. 0.2 kg
- B. 2 kg
- C. 20 kg
- D. 200 kg

**Answer:** D

**Explanation:** The total phosphorus load can be calculated as:

$$\text{Load} = \text{Concentration} \times \text{Volume}$$

Convert concentration to  $\text{kg/m}^3$ :

$$\text{Concentration} = 0.2 \text{ mg/L} = 0.0002 \text{ kg/m}^3$$

Thus,

$$\text{Load} = 0.0002 \text{ kg/m}^3 \times 1,000,000 \text{ m}^3 = 200 \text{ kg}$$

**Question 390:**

A groundwater model indicates that a well is experiencing a drawdown of 5 m after 12 hours of continuous pumping. If the well has a radius of 0.1 m and the aquifer has a hydraulic conductivity of 10 m/day, what is the estimated specific yield of the aquifer?

- A. 0.01
- B. 0.05
- C. 0.1
- D. 0.15

**Answer:** B

**Explanation:** The specific yield can be calculated using the relationship:

$$\text{Specific Yield} = \frac{\text{Drawdown}}{\text{Time}} \times \frac{1}{\text{Hydraulic Conductivity}}$$

Thus,

$$\text{Specific Yield} = \frac{5 \text{ m}}{12 \times 3600 \text{ s}} \times \frac{1}{10 \text{ m/day}} = 0.05$$

**Question 391:**

A hydrogeologist is evaluating a confined aquifer that has a hydraulic conductivity of 25 m/day and a thickness of 30 m. If the aquifer is being recharged at a rate of 0.1 m/year, what is the estimated sustainable yield of the aquifer over an area of 2 hectares?

- A. 2000 m<sup>3</sup>/yr
- B. 1500 m<sup>3</sup>/yr
- C. 1600 m<sup>3</sup>/yr
- D. 1700 m<sup>3</sup>/yr

**Answer:** A

**Explanation:** The sustainable yield can be estimated using:

$$\text{Sustainable Yield} = \text{Recharge Rate} \times \text{Area}$$

Convert the recharge rate to meters:

$$\text{Recharge Rate} = 0.1 \text{ m/yr}$$

Convert area to square meters:

$$\text{Area} = 2 \text{ hectares} = 20,000 \text{ m}^2$$

Thus,

$$\text{Sustainable Yield} = 0.1 \text{ m/yr} \times 20,000 \text{ m}^2 = 2,000 \text{ m}^3/\text{yr}$$

**Question 392:**

An engineer is analyzing groundwater flow through a heterogeneous aquifer. The hydraulic gradient in one section of the aquifer is measured at 0.03, and the hydraulic conductivity is 12 m/day. What is the groundwater flow velocity in that section?

- A. 0.36 m/day
- B. 0.48 m/day
- C. 0.56 m/day
- D. 0.72 m/day

**Answer:** A

**Explanation:** Groundwater flow velocity can be calculated using Darcy's law:

$$v = K \cdot i$$

Where  $K$  is hydraulic conductivity and  $i$  is hydraulic gradient.

Thus,

$$v = 12 \text{ m/day} \times 0.03 = 0.36 \text{ m/day}$$

**Question 393:**

A well in an unconfined aquifer is pumped at a rate of 100 L/s. After 48 hours of continuous pumping, the water level in the well has dropped from 15 m to 10 m. What is the total drawdown experienced by the well?

- A. 2 m
- B. 3 m
- C. 4 m
- D. 5 m

**Answer:** D

**Explanation:** The drawdown is calculated as:

$$\text{Drawdown} = \text{Initial Water Level} - \text{Final Water Level}$$

Thus,

$$\text{Drawdown} = 15 \text{ m} - 10 \text{ m} = 5 \text{ m}$$

**Question 394:**

A civil engineer is studying the impact of a wastewater discharge on a river's dissolved oxygen (DO) levels. If the river has a flow rate of  $4 \text{ m}^3/\text{s}$  and the DO concentration downstream of the discharge is  $5 \text{ mg/L}$ , while the upstream concentration is  $8 \text{ mg/L}$ , what is the total mass of oxygen depleted over a 24-hour period?

- A. 1288 kg
- B. 1576 kg
- C. 1036 kg
- D. 1296 kg

**Answer:** C

**Explanation:** The mass of oxygen lost can be calculated as:

$$\text{Mass Loss} = (\text{Upstream DO} - \text{Downstream DO}) \times \text{Flow Rate} \times \text{Time}$$

Where:

$$\text{Mass Loss} = (8 \text{ mg/L} - 5 \text{ mg/L}) \times 4 \text{ m}^3/\text{s} \times 86,400 \text{ s}$$

Convert mg/L to kg/m<sup>3</sup>:

$$\text{Mass Loss} = 3 \text{ mg/L} \times 4 \times 86,400 = 1036.8 \text{ kg}$$

**Question 395:**

An environmental scientist is calculating the Total Maximum Daily Load (TMDL) for nitrogen in a river. The current nitrogen load is 2,200 kg/year, and the TMDL is set at 1,500 kg/year. What is the percentage reduction needed to meet the TMDL?

- A. 25%
- B. 32%
- C. 40%
- D. 50%

**Answer:** B

**Explanation:** The percentage reduction can be calculated as:

$$\text{Reduction} = \frac{\text{Current Load} - \text{TMDL}}{\text{Current Load}} \times 100$$

Thus,

$$\text{Reduction} = \frac{2200 - 1500}{2200} \times 100 \approx 31.82\%$$

**Question 396:**

A lake has a total phosphorus concentration of 0.15 mg/L. If the lake has a

volume of 500,000 m<sup>3</sup>, what is the total phosphorus load in kilograms?

- A. 10.75 kg
- B. 11.25 kg
- C. 15.00 kg
- D. 75.0 kg

**Answer:** D

**Explanation:** The total phosphorus load can be calculated as:

$$\text{Load} = \text{Concentration} \times \text{Volume}$$

Convert concentration to kg/m<sup>3</sup>:

$$\text{Load} = 0.15 \text{ mg/L} \times 500,000 \text{ m}^3 = 75 \text{ kg}$$

**Question 397:**

In a groundwater contamination study, a monitoring well shows a concentration of benzene at 5 µg/L. If the well extracts water at a rate of 10 L/min, what is the total mass of benzene extracted in a 30-minute sampling period?

- A. 0.15 mg
- B. 0.25 mg
- C. 0.50 mg
- D. 1.50 mg

**Answer:** D

**Explanation:** The total mass can be calculated as:

$$\text{Mass} = \text{Concentration} \times \text{Flow Rate} \times \text{Time}$$

Convert flow rate to L/h:

$$\text{Mass} = 5 \mu\text{g}/\text{L} \times 10 \text{ L}/\text{min} \times 30 \text{ min} = 1,500 \mu\text{g} = 1.5 \text{ mg}$$

**Question 398:**

A civil engineer is evaluating a stream's health by assessing its biological oxygen demand (BOD). If the natural BOD of the stream is 4 mg/L and the BOD after a pollutant influx is measured at 12 mg/L, what is the increase in BOD due to the pollutants?

- A. 4 mg/L
- B. 6 mg/L
- C. 8 mg/L
- D. 10 mg/L

**Answer:** C

**Explanation:** The increase in BOD is calculated as:

$$\text{Increase in BOD} = \text{Post-Pollution BOD} - \text{Natural BOD}$$

Thus,

$$\text{Increase in BOD} = 12 \text{ mg}/\text{L} - 4 \text{ mg}/\text{L} = 8 \text{ mg}/\text{L}$$

**Question 399:**

A groundwater model reveals that a well has a drawdown of 3 m after 24 hours of pumping at a rate of 80 L/s. If the well has a radius of 0.15 m, what is the specific capacity of the well in L/s/m?

- A. 15.33 L/s/m
- B. 26.67 L/s/m
- C. 10.00 L/s/m
- D. 12.00 L/s/m

**Answer:** B

**Explanation:** Specific capacity can be calculated using:

$$\text{Specific Capacity} = \frac{\text{Discharge Rate}}{\text{Drawdown}}$$

Thus,

$$\text{Specific Capacity} = \frac{80 \text{ L/s}}{3 \text{ m}} \approx 26.67 \text{ L/s/m}$$

**Question 400:**

An environmental engineer is assessing the impact of nutrient runoff on a pond. If the pond has a surface area of 1 hectare and receives 15 kg of phosphorus from runoff annually, what is the concentration of phosphorus in mg/L, assuming an average depth of 2 m?

- A. 0.15 mg/L
- B. 0.75 mg/L
- C. 91.00 mg/L
- D. 750 mg/L

**Answer:** D

**Explanation:** Convert area to square meters:

$$\text{Area} = 1 \text{ hectare} = 10,000 \text{ m}^2$$

The volume of the pond is:

$$\text{Volume} = \text{Area} \times \text{Depth} = 10,000 \text{ m}^2 \times 2 \text{ m} = 20,000 \text{ m}^3$$

Convert kg to mg:

$$\text{Concentration} = \frac{15 \text{ kg} \times 1,000,000 \text{ mg/kg}}{20,000 \text{ m}^3} = 750 \text{ mg/L}$$

**Question 401:**

A stream has a flow rate of  $1.5 \text{ m}^3/\text{s}$  and a dissolved oxygen (DO) concentration of  $9 \text{ mg/L}$  upstream. If the DO concentration drops to  $5 \text{ mg/L}$  downstream after discharge from a wastewater treatment plant, what is the total mass of oxygen lost in kilograms over 24 hours?

- A. 518.4 kg
- B. 288 kg
- C. 864 kg
- D. 1,728 kg

**Answer:** A

**Explanation:** The mass of oxygen lost can be calculated as:

$$\text{Mass Loss} = (\text{Upstream DO} - \text{Downstream DO}) \times \text{Flow Rate} \times \text{Time}$$

Thus,

$$\text{Mass Loss} = (9 \text{ mg/L} - 5 \text{ mg/L}) \times 1.5 \text{ m}^3/\text{s} \times 86,400 \text{ s}$$

Convert  $\text{mg/L}$  to  $\text{kg/m}^3$ :

$$\text{Mass Loss} = 4 \text{ mg/L} \times 1.5 \text{ m}^3/\text{s} \times 86,400 \text{ s} = 518,400 \text{ mg} = 518.4 \text{ kg}$$

**Question 402:**

In a water quality assessment, a river's total nitrogen concentration is measured at  $12 \text{ mg/L}$ . If the river has a flow rate of  $2.5 \text{ m}^3/\text{s}$ , what is the total nitrogen load in kilograms per day?

- A. 1250 kg/day
- B. 1300 kg/day
- C. 1036 kg/day
- D. 1600 kg/day

**Answer:** C

**Explanation:** The nitrogen load can be calculated as:

$$\text{Load} = \text{Concentration} \times \text{Flow Rate} \times \text{Time}$$

Thus,

$$\text{Load} = 12 \text{ mg/L} \times 2.5 \text{ m}^3/\text{s} \times 86,400 \text{ s} = 1,036,800 \text{ mg} = 1,036.8 \text{ kg}$$



# KILLEXAMS.COM

Killexams.com is an online platform that offers a wide range of services related to certification exam preparation. The platform provides actual questions, exam dumps, and practice tests to help individuals prepare for various certification exams with confidence. Here are some key features and services offered by Killexams.com:



**Actual Exam Questions:** Killexams.com provides actual exam questions that are experienced in test centers. These questions are updated regularly to ensure they are up-to-date and relevant to the latest exam syllabus. By studying these actual questions, candidates can familiarize themselves with the content and format of the real exam.

**Exam Dumps:** Killexams.com offers exam dumps in PDF format. These dumps contain a comprehensive collection of questions and answers that cover the exam topics. By using these dumps, candidates can enhance their knowledge and improve their chances of success in the certification exam.

**Practice Tests:** Killexams.com provides practice tests through their desktop VCE exam simulator and online test engine. These practice tests simulate the real exam environment and help candidates assess their readiness for the actual exam. The practice tests cover a wide range of questions and enable candidates to identify their strengths and weaknesses.

**Guaranteed Success:** Killexams.com offers a success guarantee with their exam dumps. They claim that by using their materials, candidates will pass their exams on the first attempt or they will refund the purchase price. This guarantee provides assurance and confidence to individuals preparing for certification exams.

**Updated Content:** Killexams.com regularly updates its question bank and exam dumps to ensure that they are current and reflect the latest changes in the exam syllabus. This helps candidates stay up-to-date with the exam content and increases their chances of success.

**Technical Support:** Killexams.com provides free 24x7 technical support to assist candidates with any queries or issues they may encounter while using their services. Their certified experts are available to provide guidance and help candidates throughout their exam preparation journey.